



King's Research Portal

DOI:

[10.1016/j.ophtha.2018.03.026](https://doi.org/10.1016/j.ophtha.2018.03.026)

Document Version

Peer reviewed version

[Link to publication record in King's Research Portal](#)

Citation for published version (APA):

European Eye Epidemiology (E3) Consortium, Mauschwitz, M. M., Bonnemaier, P. W. M., Diers, K., Rauscher, F. G., Elze, T., Engel, C., Loeffler, M., Colijn, J. M., Ikram, M. A., Vingerling, J. R., Williams, K. M., Hammond, C. J., Creuzot-Garcher, C., Bron, A. M., Silva, R., Nunes, S., Delcourt, C., Cougnard-Grégoire, A., ... Finger, R. P. (2018). Systemic and Ocular Determinants of Peripapillary Retinal Nerve Fiber Layer Thickness Measurements in the European Eye Epidemiology (E3) Population. *Ophthalmology*.
<https://doi.org/10.1016/j.ophtha.2018.03.026>

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

Systemic and ocular determinants of peripapillary retinal nerve fiber layer thickness measurements in the E3 population

Matthias M. Mauschitz, MD^{1,2}, Pieter W. M. Bonnemaier, MD^{3,4}, Kersten Diers, MSc¹, Franziska G. Rauscher, PhD^{5,7}, Tobias Elze, PhD^{5,6}, Christoph Engel, MD, PhD^{5,7}, Markus Loeffler, MD, PhD^{5,7}, Johanna Maria Colijn, MD, MSc^{3,4}, M. Arfan Ikram, MD, PhD⁴, Johannes R. Vingerling, MD, PhD³, Katie M. Williams, MD, PhD⁸, Christopher J. Hammond, MD, PhD⁸, Catherine Creuzot-Garcher, MD, PhD^{9,10}, Alain M. Bron, MD, PhD^{9,10}, Rufino Silva, MD, PhD^{11,12,13}, Sandrina Nunes, PhD¹³, Cécile Delcourt, PhD¹⁴, Audrey Cougnard-Grégoire, PhD¹⁴, Frank G. Holz, MD², Caroline C. W. Klaver, MD, PhD^{3,4}, Monique M. B. Breteler, MD, PhD^{1,15}, Robert P. Finger, MD, PhD²

On behalf of the European Eye Epidemiology (E3) Consortium

¹Population Health Sciences, German Centre for Neurodegenerative Diseases (DZNE), Bonn, Germany

²Department of Ophthalmology, University of Bonn, Germany

³Department of Ophthalmology, Erasmus MC, Rotterdam, the Netherlands

⁴Department of Epidemiology, Erasmus MC, Rotterdam, the Netherlands

⁵Leipzig Research Centre for Civilization Diseases, Leipzig University, Leipzig, Germany

⁶Schepens Eye Research Institute, Harvard Medical School, Boston, MA, USA

⁷Institute for Medical Informatics, Statistics and Epidemiology, Leipzig University, Leipzig, Germany

⁸Section of Academic Ophthalmology, School of Life Course Sciences, FoLSM, King's College London, London, UK

⁹Department of Ophthalmology, University Hospital Dijon

¹⁰Eye and nutrition research group, University of Bourgogne Franche Comté, France

¹¹Department of Ophthalmology, Centro Hospitalar e Universitário de Coimbra (CHUC), Portugal

¹²Faculty of Medicine, University of Coimbra, Institute for Biomedical Imaging and Life Sciences (FMUC-IBILI), Portugal

¹³Association for Innovation and Biomedical Research on Light and Image (AIBILI), Coimbra, Portugal

¹⁴University of Bordeaux, Inserm, Bordeaux Population Health Research Center, Team LEHA, UMR 1219, F-33000 Bordeaux, France.

¹⁵Institute for Medical Biometry, Informatics and Epidemiology, University of Bonn, Faculty of Medicine, Germany

Running head: Determinants of pRNFLT in the E3 - population

Corresponding author:

Robert Patrick Finger

Department of Ophthalmology

University of Bonn, Germany

Ernst-Abbe-Straße 2, 53127 Bonn, Germany

Email: robert.finger@ukbonn.de

+49 228 287 11764

Funding:

The Alienor Study was supported by Laboratoires Théa (Clermont-Ferrand, France); University of Bordeaux (Bordeaux, France); Fondation Voir et Entendre (Paris, France); Caisse Nationale de Solidarité pour l'Autonomie CNSA (CNSA).

The Coimbra Study is an Investigator Initiated Study financially supported by Novartis Pharma AG.

This publication is supported by the Leipzig Research Centre for Civilization Diseases (LIFE), an organizational unit affiliated to the Medical Faculty of Leipzig University. LIFE is funded by means of the European Union, by the European Regional Development Fund (ERDF) and by funds of the Free State of Saxony within the framework of the excellence initiative (project numbers: 713-241202, 713-241202, 14505/2470, 14575/2470). Dr. Tobias Elze is supported by the following organizations and grants: BrightFocus Foundation, Lions Foundation, Grimshaw-Gudewicz Foundation, Research to Prevent Blindness, and NEI Core Grant P30EYE003790.

Montrachet Study: Funding was provided by an Inter-regional grant (PHRC) and the Regional Council of Burgundy. This study was also funded by INRA, CNRS, Université de Bourgogne, Regional Council of Burgundy France (PARI Agrale 1), FEDER (European Funding for Regional Economic Development) and French Government grant managed by the French National Research Agency (ANR) as part of the "Investissements d'Avenir" program (reference ANR-11-LABX-0021-01-LipSTIC Labex).

The Rotterdam Study is funded by Erasmus MC and Erasmus University, Rotterdam, Netherlands Organization for the Health Research and Development (ZonMw), the Research Institute for Diseases in the Elderly (RIDE), the Ministry of Education, Culture and Science, the Ministry for Health, Welfare and Sports, the European Commission (DG XII), and the Municipality of Rotterdam. SENSE-COG consortium has received funding from the European Union's Horizon 2020 research and innovation program under grant agreement No 668648. Stichting Lijf en Leven, Krimpen aan de Lek; MD Fonds, Utrecht; Rotterdamse Vereniging Blindenbelangen, Rotterdam; Stichting Oogfonds Nederland, Utrecht; Blindenpenning, Amsterdam; Blindenhulp, The Hague; Algemene Nederlandse Vereniging ter Voorkoming van Blindheid (ANVVB), Doorn; Landelijke Stichting voor Blinden en Slechtzienden, Utrecht; Swart van Essen, Rotterdam; Stichting Winckel-Sweep, Utrecht; Henkes Stichting, Rotterdam; Laméris Ootech BV, Nieuwegein; Medical Workshop, de Meern; Topcon Europe BV, Capelle aan de IJssel, all in The Netherlands, and Heidelberg Engineering, Dossenheim, Germany. Also supported by the NWO Graduate Programme 2010 BOO ([022.002.023](#); HS),

the National Institute of Health (Bethesda, MD, USA) Grants R01 EY019112 and R01 EY018853, Veterans Administration Grant I01 CX000119, and the Arnold and Mabel Beckman Initiative for Macular Research. TwinsUK phenotyping was funded by the International Glaucoma Association (2013 research award) and the Wellcome Trust. The study also receives support from the National Institute for Health Research (NIHR)-funded BioResource, Clinical Research Facility and Biomedical Research Centre based at Guy's and St Thomas' NHS Foundation Trust in partnership with King's College London. Dr. Robert Finger's research group receives funding from the Else Kröner-Fresenius-Stiftung (GSO/EKFS 16) and the Jackstädt Stiftung.

Conflict of interest: Alain M. Bron reports personal fees from Allergan, personal fees from Bausch Lomb, grants from Horus, personal fees from Théa, personal fees from Carl Zeiss Meditec, outside the submitted work. Catherine Creuzot-Garcher reports grants and personal fees from Allergan, personal fees from Bayer, personal fees and other from Novartis, grants from Horus, grants and personal fees from Thea, outside the submitted work. Cécile Delcourt is consultant for Allergan, Bausch+Lomb, Laboratoires Théa, Novartis and Roche, outside the submitted work. Frank G. Holz reports fees from Heidelberg Engineering, Optos, Zeiss, Genentech, Acucela, Bayer Healthcare and Novartis, outside the submitted work. Robert P. Finger reports fees from Novartis, Bayer, Abbvie, Opthea, Novelion, Retinal Implant and Santen, outside the submitted work. Rufino Silva is member of Advisory Board for Novartis, Bayer, Allergan, Alimera, Alcon, THEA.

Abstract

Objective: To investigate systemic and ocular determinants of peripapillary retinal nerve fiber layer thickness (pRNFLT) in the European population.

Design: Cross-sectional meta-analysis.

Participants: 16,084 European adults from eight cohort studies (mean age range from 56.9 ± 12.3 to 82.1 ± 4.2 years) of the European Eye Epidemiology (E3) consortium.

Methods: We examined associations with pRNFLT measured by spectral – domain optical coherence tomography in each study using multivariable linear regression and pooled results using random effects meta-analysis.

Main Outcome Measures: Determinants of pRNFLT.

Results: Mean pRNFLT ranged from 86.8 ± 21.4 in the Rotterdam Study I to 104.7 ± 12.5 μm in the Rotterdam Study III. We found the following factors to be associated with reduced pRNFLT: Older age ($\beta = -0.38$ $\mu\text{m}/\text{year}$, 95% confidence interval (CI) = $-0.57, -0.18$), higher intraocular pressure (IOP; $\beta = -0.36$ $\mu\text{m}/\text{mmHg}$, 95% CI = $-0.56, -0.15$), visual impairment ($\beta = -5.50$ μm , 95% CI = $-9.37, -1.64$) and history of systemic hypertension ($\beta = -0.54$ μm , 95% CI = $-1.01, -0.07$) and stroke ($\beta = -1.94$ μm , 95% CI = $-3.17, -0.72$). A suggestive, albeit non-significant, association was observed for dementia ($\beta = -3.11$ μm , 95% CI = $-6.22, 0.01$). Higher pRNFLT was associated with more hyperopic spherical equivalent (SE; $\beta = 1.39$ $\mu\text{m}/\text{diopter}$, 95% CI = $1.19, 1.59$) and smoking ($\beta = 1.53$ μm , 95% CI = $1.00, 2.06$ for current smokers compared to never-smokers).

Conclusions: In addition to previously described determinants such as age and refraction, we found that systemic vascular and neurovascular diseases were associated with reduced pRNFLT. These may be of clinical relevance, especially in glaucoma monitoring of patients with newly occurring vascular co-morbidities.

INTRODUCTION

The assessment of peripapillary retinal nerve fiber layer thickness (pRNFLT) with Spectral – Domain Optical Coherence Tomography (SD-OCT) has become of increasing importance in the evaluation of glaucoma and its progression^{1,2}. Although debated, pRNFLT measurements hold promise as a biomarker for neurodegenerative diseases such as Alzheimer’s disease (AD) and multiple sclerosis (MS)^{3,4}.

While pRNFLT measurements have increased insight into the development of diseases, it has been difficult to evaluate which changes fall within the physiological range. Most OCT devices compare pRNFLT measurements against reference databases that are built into the machine analysis software. These data are mostly derived from relatively small sample populations. Whether these databases adequately capture normal anatomical variation across a wide age range remains unclear.

Only few studies investigated ocular and systemic determinants of pRNFLT in the general population⁵. They reported inconsistent results for many ocular and systemic parameters including sex or body-mass-index (BMI)^{5,6}. To date, only age^{7,8}, refraction⁹ or axial length (AL)¹⁰ have been consistently associated with measured pRNFLT across studies. In addition, the majority of large-scale studies assessing these associations were performed in (young) Asian populations^{6,11–14}. It is unclear whether or not these results can be applied to European, i.e. mostly Caucasian, populations.

The purpose of this study was to assess systemic and ocular determinants of pRNFLT using pooled data from eight European population-based studies.

METHODS

Included studies

The European Eye Epidemiology (E3) consortium is a collaborative network of population-based studies across Europe with the overarching aim of developing and analyzing large pooled datasets to increase understanding of eye disease and vision loss¹⁵. For this study, we analyzed data on pRNFLT from eight different studies. The included data were cross-

sectional and the right eye was chosen to be the study eye. All studies adhered to the tenets of the Declaration of Helsinki and had local ethical committee approval. All participants gave written informed consent.

Assessments and data analyses

Retinal nerve fiber layer thickness was measured as global pRNFLT with different OCT devices, scan modalities (mostly circular scans) and automated segmentation algorithms in the respective studies (see Table 1). pRNFLT outliers were excluded prior to analyses according to Chauvenet's criterion. Briefly, depending on sample size we excluded participants with pRNFLT above or below a certain range of standard deviations from the mean¹⁶. To investigate determinants of pRNFLT, multivariable linear regression models including the variables of interest were conducted. Factors to be tested for association with pRNFLT were considered in multiple steps. As first and most important step, variables were chosen a priori based on literature and availability in the individual studies. Subsequently, we performed univariable linear regression models of potential factors at study level to assess possible impact on pRNFLT. In the last step the factors of the multivariable models were decided on as a trade-off between priority of the respective factors and the maximum possible population size of the model.

The independent variables of the multivariable linear regression model were age, sex, body-mass-index (BMI), visual impairment as defined by the World Health Organization (WHO) (best corrected visual acuity (BCVA) <0.3 decimal), intraocular pressure (IOP), spherical equivalent (SE), smoking status and history of systemic hypertension, diabetes, stroke and dementia. The multivariable regression model was conducted for each individual study and residuals were then plotted and normal distribution assessed. Since OCT devices were changed within the course of the Rotterdam Study (From 3D-OCT 1000 to 3D-OCT 2000, Topcon Medical Systems, Oakland, NJ, USA), we controlled for the OCT device in the multivariable regression models of the Rotterdam Study II and III. In the TwinsUK Study, we

performed a hierarchical multivariable regression model to control for family dependencies between twins.

Subsequently, random-effects meta-analysis was used to combine effect estimates (beta coefficients) of each individual predictor from the multivariable regression model among studies. A random-effects approach was chosen a priori based on the heterogeneity in the data caused by the different OCT devices¹⁷ and the set-up of the studies. Our analyses were conducted twice, with and without known glaucoma patients.

Not all independent variables of the multivariable regression model were available in every participating study. The multivariable regression models in the respective studies were therefore performed without the missing variables and the study was excluded from the meta-analysis of that respective missing covariate. All analyses were performed with the statistical software RStudio (R version 3.4.1, RStudio Inc., Boston, MA, <https://www.rstudio.com/>), statistical significance was set at $p < 0.05$.

RESULTS

A total of 16,084 participants from eight population-based studies were included, about one percent pRNFLT outliers per study were excluded (supplemental Table 1b). The mean age of participants ranged from 56.9 ± 12.3 years in the LIFE Study to 82.1 ± 4.2 years in the Alienor Study. Mean global pRNFLT ranged from 86.8 ± 21.4 microns in the Rotterdam Study I to 104.7 ± 12.5 microns in the Rotterdam Study III (Table 1). Further participant characteristics for each study are presented in supplemental Table 1b. The results of the multivariable regression models for each individual study are reported in Table 2. Data on dementia were only available in the Rotterdam Study cohorts and the Alienor Study. Furthermore, in the TwinsUK Study no sufficient data were available on visual impairment, glaucoma, hypertension and smoking status; in the LIFE Study, no data were available on visual impairment, SE and IOP.

In the meta-analyzed multivariable regression model (Table 3 and Figures 1a and 1b), age and IOP were negatively associated with pRNFLT, even after excluding glaucoma patients. A

history of stroke and hypertension were both associated with a reduced pRNFLT. When substituting hypertension with mean systolic blood pressure (in mmHg), no association was found.

A suggestive, but non-significant association with reduced pRNFLT was observed for dementia. Visual impairment as defined by the WHO was associated with reduced pRNFLT in the meta-analysis. We found this association in the Alienor and Rotterdam Study I-III, while there was no association in the Montrachet and Coimbra Study.

Women had a thicker pRNFLT than men in the meta-analysis. However, when correcting for AL rather than SE in the five studies with data on AL, this association disappeared. SE was positively associated with pRNFLT, even after excluding highly myopic (< -6 diopters) and highly hyperopic eyes ($> +4$ diopters) as well as eyes with pseudophakia (supplemental Figures A and B). Longer AL was associated with reduced pRNFLT in our sensitivity analyses (beta= $-3.48\mu\text{m}$ per mm longer AL, 95% CI= $-4.18, -2.77$) (supplemental Figure C). Both, former and current smoking were associated with thicker pRNFLT, but prevalence and associations differed considerably between studies. To assess the influence of education on smoking, we corrected for education and the associations persisted. After excluding data from the LIFE Study, which is the largest study with the highest proportion of smokers (data weighted $>60\%$ in the meta-analysis), the association remained significant for current but not for former smoking (supplemental Figures D-G). For BMI, we found a small but significant association with increased pRNFLT after excluding glaucoma patients. All associations except for former smoking held true after excluding the 619 known glaucoma patients (Table 3). Furthermore, we detected no relevant changes of associations when performing the multivariable regression analyses stratified by sex or when excluding the LIFE study cohort being the largest single study (results not reported).

DISCUSSION

Our study confirms the previously reported associations of age and SE with pRNFLT and identifies several additional factors associated with pRNFLT, namely IOP (even in individuals

without a history of glaucoma), stroke, hypertension and smoking. Furthermore, we found a trend of reduced pRNFLT in participants with dementia. Our results suggest that a number of ocular as well as systemic factors need to be considered when assessing pRNFLT. To date, none of this has for example been implemented as potentially influencing factors in reference databases for OCT devices or any algorithms assessing pRNFLT change.

First publications on determinants of OCT – based pRNFLT measurements reported older age and greater AL to be associated with thinner pRNFLT^{18,19}. Budenz and coworkers investigated determinants of pRNFLT in 328 normal subjects aged 18 to 85 years using time domain – optical coherence tomography (TD-OCT) and described a decrease of 2.0 microns pRNFLT per decade and a decrease of 2.2 microns per millimeter AL¹⁹. These estimates are smaller but still compare to our results (decrease of 3.8 microns pRNFLT on average per decade and 3.48 microns per millimeter AL). A subsequent study evaluated determinants of pRNFLT in 542 healthy adults aged 40 to 80 years using SD – OCT (Cirrus HD-OCT; Carl Zeiss Meditec, Inc., Dublin, CA) and confirmed the associations of pRNFLT with age and AL¹¹.

Subsequently, larger population studies mostly from Asia were conducted to investigate further determinants of pRNFLT. We have affirmed results from the Beijing Eye Study in 2548 participants considering the influence of age and refractive error. That study also showed a higher pRNFLT of 2.9 microns in women¹⁴, in keeping with our results of women having a higher pRNFLT of 2.2 microns. Similar to our models, the Beijing Eye Study corrected for refractive error instead of actual AL. Interestingly, after correcting for AL in our analyses, sex was no longer associated with pRNFLT. Based on this, we hypothesize that AL, which is on average shorter in women, confounds the effect of sex on pRNFLT. In general, SE is a good proxy for AL and we found a strong association of higher SE with thicker pRNFLT, even in both our sensitivity analyses, which eliminated subjects with high refractive errors. The underlying mechanisms of the association of longer AL and thinner pRNFLT are arguable²⁰. Frequently suggested mechanisms are either a stretching due to a longer eye bulb or artificially decreased measurements due to magnification^{21,22}. However,

irrespective of the causal mechanism, the clinical relevance of adjusting for refraction or AL in OCT – imaging seems obvious.

Higher IOP was associated with reduced pRNFLT in our analyses even after excluding known glaucoma patients. However, since glaucoma was self-reported in some of the participating studies, not all actual glaucoma patients might have been excluded in our analyses. Visual impairment (BCVA < 0.3 decimal) as a proxy for any ocular pathology was associated with thinner pRNFLT in the Alienor Study and all of the Rotterdam Studies. The Coimbra and Montrachet Study were likely underpowered to find an effect, because of very few cases with reduced BCVA in these studies.

Previous studies reported contradictory results on the impact of hypertension and blood pressure on pRNFLT^{9,23,24}. Our results show reduced pRNFLT in hypertensive patients, but no association of pRNFLT with actual systolic blood pressure. Blood pressure measurements, however, are known to vary with method and associations with systolic blood pressure may have been masked by any use of antihypertensive medication. In contrast to hypertension, most studies investigating the effect of diabetes on pRNFLT report diabetic patients to have thinner pRNFLT^{25,26}. This is in not agreement with our results that do not show an association of reduced pRNFLT in diabetic patients. Nether the less, we hypothesize that microvascular pathology and ischemia due to hypertension and/or diabetes may be a cause for reduced pRNFLT, as it has been suggested previously²⁵.

Both, former and current smoking were associated with thicker pRNFLT in our meta-analysis, even in several sensitivity analyses including correction for educational level. This association does not seem biologically plausible given the observed pRNFLT decrease in metabolic diseases. Potential biologic explanations could be reduced axonal flow or axonal swelling in the course of axonal degeneration due to intake of neurotoxins and cytotoxins from cigarette smoke. However, our results are in contrast with findings of earlier studies^{27,28}, which reported reduced pRNFLT in smokers. Suggested mechanisms leading to decreased pRNFLT were toxic damage through free radicals, increased IOP and reduced perfusion^{27–29}. We controlled for IOP as well as hypertension and diabetes, which all may influence

perfusion. It is therefore unclear what might explain this association. Current smokers were on average younger in our participating studies compared to never and former smokers. Hence, even though we controlled for age in our models, we cannot entirely rule out residual confounding. Additionally, the E3 studies are not representative studies of European populations and smoking percentages therefore do not reflect actual percentages. There was heterogeneity between studies considering smoking prevalence and oppositional effects of former smoking in some studies. After excluding the LIFE Study, which was dominantly weighted in the smoking meta-analysis, the Rotterdam Study III showed to be weighted strongest for current smoking. When excluding also the Rotterdam Study III, the impact of smoking is weakened but holds true. Still, the associations seem to be particularly driven by the large studies. This is also underlined by increasing heterogeneity for former and current smoking in the meta-analysis after excluding the LIFE Study. Moreover, there is no information on the time interval between cessation of smoking and OCT – imaging for the former smokers, which may have an impact, as well. Further studies are needed to confirm or refute our observation, which may well be a chance finding.

Past studies have reported stroke patients to have thinner pRNFLT, which was hypothesized to be caused by transneuronal retrograde degeneration^{30,31}. Our data confirm the association of stroke and decreased pRNFLT. Additionally, in dementia patients we found a trend of reduced pRNFLT. Again, this is in accordance to various previous studies, which report dementia patients to have reduced pRNFLT^{4,32}. Thus far, the underlying mechanisms remain unclear. Loss of peripapillary RNFL is a hallmark of glaucoma and longitudinal pRNFLT evaluation is a crucial part of glaucoma management. In our meta-analysis, all associations persisted after excluding known glaucoma patients except for former smoking. This indicates that the detected determinants are independent of the presence of glaucoma.

As described previously, structural decline of pRNFLT occurs before functional loss in perimetry in glaucoma patients. An earlier study reported the difference in pRNFLT between glaucomatous and healthy eyes eight years before the onset of visual field impairment to be around 5 μm ³³. This is in the range of some associations found in our study and underlines

the potential impact on the interpretation of pRNFLT. Our results have two main clinical implications. Firstly, the normative databases built into the devices should reflect our results, when presenting normal values for pRNFLT. Also, presence of vascular disease including a history of stroke should be considered when defining normative datasets or when clinically evaluating pRNFLT. As discussed above, the magnitude of impact of the respective determinants may have clinical relevance, especially in the presence of more than one factor reducing pRNFLT. Secondly, in glaucoma or other patients followed up with pRNFLT measurements, an incident stroke or dementia may cause a decrease in pRNFLT, which would not primarily be due to glaucoma or other ocular disease progression. For example, this may simulate an aggravation of glaucoma and needs to be considered by the clinician when tailoring the glaucoma management.

The strengths of this study consist of the large pooled sample combining data of eight studies from five European countries. To our knowledge, this study represents the largest European study on determinants of pRNFLT thus far. As mentioned, previous population studies reporting data on associations with pRNFLT were conducted in mostly Asian populations and results cannot directly be transferred to European individuals. The associations of this study were assessed in meta-analyses of all participating populations, thus they are not limited to one single population only. This reduces the possibility that an association was solely due to chance within one population and increases generalizability. However, several limitations of our study need to be considered. The use of different OCT-devices between studies may have increased variability and prohibited direct pooling of pRNFLT data. To overcome this lack of direct comparability we performed the analysis separately within studies and then pooled studies' effect estimates using random-effect meta-analysis. Furthermore, we found no interactions between type of device and any predictor variable in additional sensitivity analyses in the Rotterdam Study II and III, which had a device upgrade within course of the study. However, residual influence of different OCT devices cannot be entirely excluded. As expected when combining different large-scale population studies, we observed between study heterogeneity for the independent variables

and their influence on pRNFLT. The degree of heterogeneity of the respective covariates was assessed using the I² – statistics and ranged from 0% to 97% (see Table 3). As described, this heterogeneity between studies was addressed by using random effect meta-analysis¹⁷. In accordance with previous literature, the relationship between pRNFLT and age was linear in our sample. Having no data for children and young adults, we do not know whether the relationship between pRNFLT and age is strictly linear throughout life but would assume so based on our data. Thus, we investigated associations using multivariable linear regression modeling. Based on this, any non-linear relationships may have been underrepresented. Quality control was performed within each study differently (supplemental Table 2). Some studies performed manual (re)-segmentation, excluded OCT images below a certain scan quality and scans with artifacts, while others included all scans with sufficient quality as evaluated by the performing technician. As sensitivity analysis we excluded participants with an image quality value below 45 (as recommended by the manufacturer) in the Rotterdam Studies I-III. We found no relevant changes of direction in any association, but the confidence intervals became broader due to a reduced sample size (supplemental Table 3). Hence, even though the lack of centralized quality control is a limitation to our analyses, the impact of poor quality scans seems to be low as indicated by our supplemental sensitivity analyses. Within each study, the number of participants in which OCT imaging could not be performed or in which the images were of low quality and thus unusable is a small proportion only (supplemental Table 2). For example, in the Rotterdam Study I-III the number of participants with no or insufficient OCT data was 10%, 6% and 15%, respectively. These subjects were older and more likely to have stroke (RS I), dementia (RS II and III) and hypertension (RS III) than the included participants. This indicates that our associations may be underestimations of the true effect. Several independent variables were not available in some studies. Therefore not all multivariable models could be corrected for all variables. However, no relevant differences of associations were detectable, when comparing studies with and studies without any missing data. Hence, the absence of certain variables in some studies did not relevantly alter the associations of the available data. Methods of

assessments varied between our studies. This concerns e.g. the best-corrected visual acuity, which was sometimes measured subjectively and sometimes by autorefractor. In addition, information on diseases was assessed differently. While glaucoma was defined based on optic disc evaluation and perimetry in the Alienor Study and Rotterdam Study I-III, it was self-reported in the LIFE Study. Furthermore, we did not distinguish between the various types of dementia, which may have different impact on pRNFLT. These differences contribute again to larger heterogeneity and the relation between self-reported diseases and pRNFLT may have been estimated with less precision. Lastly, our data were cross-sectional only, thus causal deductions from the detected associations are limited and further longitudinal studies are needed.

In conclusion, the current analyses identified important additional determinants of pRNFLT, which should be considered when assessing pRNFLT both clinically and in epidemiological research. The magnitude of changes in pRNFLT by determinant is likely clinically relevant and the biology of pRNFLT thinning is complex, with mechanical pressure, microvascular ischemia and neuronal degeneration being implied. This is reflected in the complexity of factors, which influence pRNFLT and hence need to be considered. In particular, the associations with systemic vascular and neurovascular diseases merit further research.

Acknowledgments

The authors would like to gratefully acknowledge the contribution of the following persons: Alberta A. H. J. Thiadens (Rotterdam), Nomdo M. Jansonius (Groningen/Rotterdam) and Paulus de Jong (Rotterdam). The authors would additionally like to thank Dr Matthias Nuechter (LIFE) for his enthusiastic assistance to this collaborative research, Dr. Kerstin Wirkner (LIFE) and her team for data acquisition, and Dr. Toralf Kirsten (LIFE) for IT structure and overall support. Furthermore, the authors express sincere thanks to Verena Brendler (LIFE) and Yvonne Dietz (LIFE) for data management.

Members of the European Eye Epidemiology (E3) Consortium:

Last name	First name	Institution	City	Country
Acar	Niyazi	Inra-University of Burgundy	Dijon	France
Anastosopoulos	Eleftherios	University of Thessaloniki	Thessaloniki	Greece
Azuara-Blanco	Augusto	Queen's University	Belfast	UK
Berendschot	Tos	University Eye Clinic Maastricht	Maastricht	Netherlands
Berendschot	Tos	University of Maastricht	Maastricht	Netherlands
Bergen	Arthur	Netherlands Institute for Neurosciences-KNAW	Amsterdam	Netherlands
Bertelsen	Geir	University of Tromso	Tromso	Norway
Biquet	Christine	University Hospital of Dijon	Dijon	France
Bird	Alan	Moorfield's Eye Hospital	London	UK
Bobak	Martin	Lithuanian University of health sciences	Kaunas	Lithuania
Bøgelund Larsen	Morten	University of Southern Denmark / Odense University Hospital	Odense	Denmark
Boon	Camiel	Leiden University Medical Center	Leiden	Netherlands
Bourne	Rupert	University of Ruskin	Cambridge	England
Brétilon	Lionel	Inra-University of Burgundy	Dijon	France
Broe	Rebecca	University of Southern Denmark	Odense	Denmark
Bron	Alain	University Hospital of Dijon	Dijon	France
Buitendijk	Gabrielle	Erasmus Medical Center	Rotterdam	Netherlands
Cachulo	Maria Luz	AIBILI/CHUC	Coimbra	Portugal
Capuano	Vittorio	University Hospital of Créteil	Créteil	France
Carrière	Isabelle	Inserm U1061	Montpellier	France
Chakravarthy	Usha	Queen's University	Belfast	UK
Chan	Michelle	UCL Institute of Ophthalmology	London	UK
Chang	Petrus	University of Bonn	Bonn	Germany
Colijn	Johanna	Erasmus Medical Center	Rotterdam	Netherlands
Cougnard-Grégoire	Audrey	Bordeaux Population Health Research Center UMR1219	Bordeaux	France
Cree	Angela	University of Southampton	Southampton	UK
Creuzot-Garcher	Catherine	University Hospital of Dijon	Dijon	France
Cumberland	Phillippa	UCL Institute of Child Health	London	UK
Cunha-Vaz	José	AIBILI/CHUC	Coimbra	Portugal
Daen	Vincent	Inserm U1061	Montpellier	France

Mauschitz et al., Systemic and ocular determinants of peripapillary retinal nerve fiber layer thickness measurements in the E3 population

De Jong	Eiko	Radboud University	Nijmegen	Netherlands
Deak	Gabor	Medical University of Vienna	Vienna	Austria
Delcourt	Cécile	Bordeaux Population Health Research Center UMR1219	Bordeaux	France
Delyfer	Marie-Noëlle	Bordeaux Population Health Research Center UMR1219	Bordeaux	France
den Hollander	Anneke	Radboud University	Nijmegen	Netherlands
Dietzel	Martha	University of Muenster	Muenster	Germany
Erke	Maja Gran	University of Tromso	Tromso	Norway
Faria	Pedro	AIBILI/CHUC	Coimbra	Portugal
Farinha	Claudia	AIBILI/CHUC	Coimbra	Portugal
Fauser	Sascha	University Eye Hospital	Cologne	Germany
Finger	Robert	University of Bonn	Bonn	Germany
Fletcher	Astrid	London School of Hygiene and Tropical Medicine	London	UK
Foster	Paul	UCL Institute of Ophthalmology	London	UK
Founti	Panayiota	University of Thessaloniki	Thessaloniki	Greece
Gorgels	Theo	Netherlands Institute for Neurosciences-KNAW	Amsterdam	Netherlands
Grauslund	Jakob	University of Southern Denmark	Odense	Denmark
Grus	Franz	University Medical Center Mainz	Mainz	Germany
Hammond	Christopher	King's College	London	UK
Hense	Hans-Werner	University of Muenster	Muenster	Germany
Hermann	Manuel	University Eye Hospital	Cologne	Germany
Hoehn	René	University Medical Center	Mainz	Germany
Hogg	Ruth	Queen's University	Belfast	UK
Holz	Frank	University of Bonn	Bonn	Germany
Hoyng	Carel	Radboud University	Nijmegen	Netherlands
Jansonius	Nomdo	Erasmus Medical Center	Rotterdam	Netherlands
Janssen	Sarah	Netherlands Institute for Neurosciences-KNAW	Amsterdam	Netherlands
Kersten	Eveline	Radboud University	Nijmegen	Netherlands
Khawaja	Anthony	NIHR Biomedical Research Centre, Moorfields Eye Hospital NHS Foundation Trust and UCL Institute of Ophthalmology	London	UK
Klaver	Caroline	Erasmus Medical Center	Rotterdam	Netherlands
Korobelnik	Jean-François	Bordeaux Population Health Research Center UMR1219	Bordeaux	France
Lamparter	Julia	University Medical Center Mainz	Mainz	Germany
Le Goff	Mélanie	Bordeaux Population Health Research Center UMR1219	Bordeaux	France
Lechanteur	Yara	Radboud University	Nijmegen	Netherlands
Lehtimäki	Terho	Fimlab Laboratories and School of Medicine, University of Tampere	Tampere	Finland
Leung	Irene	Moorfield's Eye Hospital	London	UK
Lotery	Andrew	University of Southampton	Southampton	UK
Mauschitz	Matthias	University of Bonn	Bonn	Germany
Meester	Magda	Erasmus Medical Center	Rotterdam	Netherlands
Merle	Bénédicte	Bordeaux Population Health Research Center UMR1219	Bordeaux	France
Meyer zu Westrup	Verena	University of Muenster	Muenster	Germany
Midena	Edoardo	University of Padova	Padova	Italy
Miotto	Stefania	University of Padova	Padova	Italy

Mauschitz et al., Systemic and ocular determinants of peripapillary retinal nerve fiber layer thickness measurements in the E3 population

Mirshahi	Alireza	Dardenne Eye Hospital	Bonn	Germany
Mohan-Saïd	Sadek	Institut de la Vision	Paris	France
Mueller	Michael	Pirkanmaa Hospital District	Tampere	Finland
Muldrew	Alyson	Queen's University	Belfast	UK
Murta	Joaquim	AIBILI/CHUC	Coimbra	Portugal
Nickels	Stefan	University Medical Center	Mainz	Germany
Nunes	Sandrina	AIBILI/CHUC	Coimbra	Portugal
Owen	Christopher	University of London	London	UK
Peto	Tunde	Queen's University	Belfast	UK
Pfeiffer	Norbert	University Medical Center	Mainz	Germany
Piermarocchi	Stefano	University of Padova	Padova	Italy
Prokofyeva	Elena	Scientific Institute of Public Health (WIV-ISP)	Brussels	Belgium
Rahi	Jugnoo	UCL Institute of Ophthalmology	London	UK
Raitakari	Olli	Turku University Hospital, University of Turku	Turku	Finland
Rauscher	Franziska	Leipzig University Hospital	Leipzig	Germany
Ribeiro	Luisa	AIBILI/CHUC	Coimbra	Portugal
Rougier	Marie-Bénédicte	Bordeaux Population Health Research Center UMR1219	Bordeaux	France
Rudnicka	Alicja	University of London	London	UK
Sahel	José	Institut de la Vision	Paris	France
Salonikiou	Aggeliki	University of Thessaloniki	Thessaloniki	Greece
Sanchez	Clarisa	Radboud University	Nijmegen	Netherlands
Schmitz-Valckenberg	Steffen	University of Bonn	Bonn	Germany
Schuster	Alexander	University Medical Center	Mainz	Germany
Schweitzer	Cédric	Bordeaux Population Health Research Center UMR1219	Bordeaux	France
Segato	Tatiana	University of Padova	Padova	Italy
Shehata	Jasmin	Medical University of Vienna	Vienna	Austria
Silva	Rufino	AIBILI/CHUC	Coimbra	Portugal
Silvestri	Giuliana	Queen's University	Belfast	UK
Simader	Christian	Medical University of Vienna	Vienna	Austria
Souied	Eric	University Hospital of Créteil	Créteil	France
Speckauskas	Martynas	Lithuanian University of health sciences	Kaunas	Lithuania
Springelkamp	Henriet	Erasmus Medical Center	Rotterdam	Netherlands
Tapp	Robyn	Pirkanmaa Hospital District	Tampere	Finland
Topouzis	Fotis	University of Thessaloniki	Thessaloniki	Greece
van Leeuwen	Elisa	Erasmus Medical Center	Rotterdam	Netherlands
Verhoeven	Virginie	Erasmus Medical Center	Rotterdam	Netherlands
Verzijden	Timo	Erasmus Medical Center	Rotterdam	Netherlands
Von Hanno	Therese	University of Tromsø	Tromsø	Norway
Wiedemann	Peter	Leipzig University Hospital	Leipzig	Germany
Williams	Katie	King's College London	London	UK
Wolfram	Christian	University Medical Center	Mainz	Germany
Yip	Jennifer	UCL Institute of Ophthalmology	London	UK
Zerbib	Jennyfer	University Hospital of Créteil	Créteil	France

403

404

References

1. Mwanza J-C, Oakley JD, Budenz DL, Anderson DR. Ability of cirrus HD-OCT optic nerve head parameters to discriminate normal from glaucomatous eyes. *Ophthalmology*. 2011;118(2):241. doi:10.1016/j.ophtha.2010.06.036.
2. Gardiner SK, Fortune B, Demirel S. Localized Changes in Retinal Nerve Fiber Layer Thickness as a Predictor of Localized Functional Change in Glaucoma. *Am J Ophthalmol*. 2016;170:75-82. doi:10.1016/j.ajo.2016.07.020.
3. Marziani E, Pomati S, Ramolfo P, et al. Evaluation of retinal nerve fiber layer and ganglion cell layer thickness in Alzheimer's disease using spectral-domain optical coherence tomography. *Invest Ophthalmol Vis Sci*. 2013;54(9):5953-5958. doi:10.1167/iovs.13-12046.
4. Jones-Odeh E, Hammond CJ. How strong is the relationship between glaucoma, the retinal nerve fibre layer, and neurodegenerative diseases such as Alzheimer's disease and multiple sclerosis? *Eye (Lond)*. 2015;29(10):1270-1284. doi:10.1038/eye.2015.158.
5. Rougier M-B, Korobelnik J-F, Malet F, et al. Retinal nerve fibre layer thickness measured with SD-OCT in a population-based study of French elderly subjects: the Alienor study. *Acta Ophthalmol*. 2015;93(6):539-545. doi:10.1111/aos.12658.
6. Zhu B-D, Li S-M, Li H, et al. Retinal nerve fiber layer thickness in a population of 12-year-old children in central China measured by iVue-100 spectral-domain optical coherence tomography: the Anyang Childhood Eye Study. *Invest Ophthalmol Vis Sci*. 2013;54(13):8104-8111. doi:10.1167/iovs.13-11958.
7. Chauhan BC, Danthurebandara VM, Sharpe GP, et al. Bruch's Membrane Opening Minimum Rim Width and Retinal Nerve Fiber Layer Thickness in a Normal White Population: A Multicenter Study. *Ophthalmology*. 2015;122(9):1786-1794. doi:10.1016/j.ophtha.2015.06.001.
8. Leung CKS, Yu M, Weinreb RN, et al. Retinal nerve fiber layer imaging with spectral-domain optical coherence tomography: a prospective analysis of age-related loss. *Ophthalmology*. 2012;119(4):731-737. doi:10.1016/j.ophtha.2011.10.010.
9. Schuster AK-G, Fischer JE, Vossmerbaeumer C, Vossmerbaeumer U. Determinants of peripapillary retinal nerve fiber layer thickness regarding ocular and systemic parameters - the MIPH Eye&Health Study. *Graefes Arch Clin Exp Ophthalmol*. 2016;254(10):2011-2016. doi:10.1007/s00417-016-3422-y.
10. Yamashita T, Sakamoto T, Yoshihara N, et al. Correlations Between Retinal Nerve Fiber Layer Thickness and Axial Length, Peripapillary Retinal Tilt, Optic Disc Size, and Retinal Artery Position in Healthy Eyes. *J Glaucoma*. 2016. doi:10.1097/IJG.0000000000000550.
11. Cheung CY, Chen D, Wong TY, et al. Determinants of quantitative optic nerve measurements using spectral domain optical coherence tomography in a population-based sample of non-glaucomatous subjects. *Invest Ophthalmol Vis Sci*. 2011;52(13):9629-9635. doi:10.1167/iovs.11-7481.
12. Chen L, Huang J, Zou H, et al. Retinal nerve fiber layer thickness in normal Chinese students aged 6 to 17 years. *Invest Ophthalmol Vis Sci*. 2013;54(13):7990-7997. doi:10.1167/iovs.12-11252.
13. Kang M-T, Li S-M, Li H, et al. Peripapillary retinal nerve fibre layer thickness and its association with refractive error in Chinese children: the Anyang Childhood Eye Study. *Clin Experiment Ophthalmol*. 2016;44(8):701-709. doi:10.1111/ceo.12764.
14. Zhao L, Wang Y, Chen CX, Xu L, Jonas JB. Retinal nerve fibre layer thickness measured by Spectralis spectral-domain optical coherence tomography: The Beijing Eye Study. *Acta Ophthalmol*. 2014;92(1):e35-41. doi:10.1111/aos.12240.

15. Delcourt C, Korobelnik J-F, Buitendijk GHS, et al. Ophthalmic epidemiology in Europe: the "European Eye Epidemiology" (E3) consortium. *Eur J Epidemiol.* 2016;31(2):197-210. doi:10.1007/s10654-015-0098-2.
16. Chauvenet W. *A Manual of Spherical and Practical Astronomy*. 5th ed. Philadelphia: Lippincott Company; 1906:469-566.
17. Khawaja AP, Springelkamp H, Creuzot-Garcher C, et al. Associations with intraocular pressure across Europe: The European Eye Epidemiology (E3) Consortium. *Eur J Epidemiol.* 2016. doi:10.1007/s10654-016-0191-1.
18. Nagai-Kusuhara A, Nakamura M, Fujioka M, Tatsumi Y, Negi A. Association of retinal nerve fibre layer thickness measured by confocal scanning laser ophthalmoscopy and optical coherence tomography with disc size and axial length. *Br J Ophthalmol.* 2008;92(2):186-190. doi:10.1136/bjo.2007.127480.
19. Budenz DL, Anderson DR, Varma R, et al. Determinants of normal retinal nerve fiber layer thickness measured by Stratus OCT. *Ophthalmology.* 2007;114(6):1046-1052. doi:10.1016/j.ophtha.2006.08.046.
20. Mwanza J-C, Durbin MK, Budenz DL, et al. Profile and predictors of normal ganglion cell-inner plexiform layer thickness measured with frequency-domain optical coherence tomography. *Invest Ophthalmol Vis Sci.* 2011;52(11):7872-7879. doi:10.1167/iovs.11-7896.
21. Kang SH, Hong SW, Im SK, Lee SH, Ahn MD. Effect of myopia on the thickness of the retinal nerve fiber layer measured by Cirrus HD optical coherence tomography. *Invest Ophthalmol Vis Sci.* 2010;51(8):4075-4083. doi:10.1167/iovs.09-4737.
22. Higashide T, Ohkubo S, Hangai M, et al. Influence of Clinical Factors and Magnification Correction on Normal Thickness Profiles of Macular Retinal Layers Using Optical Coherence Tomography. *PLoS ONE.* 2016;11(1):e0147782. doi:10.1371/journal.pone.0147782.
23. Sahin OZ, Sahin SB, Ayaz T, et al. The impact of hypertension on retinal nerve fiber layer thickness and its association with carotid intima media thickness. *Blood Press.* 2015;24(3):178-184. doi:10.3109/08037051.2014.1000562.
24. Topouzis F, Coleman AL, Harris A, et al. Association of blood pressure status with the optic disk structure in non-glaucoma subjects: the Thessaloniki eye study. *Am J Ophthalmol.* 2006;142(1):60-67. doi:10.1016/j.ajo.2006.02.055.
25. Jeon SJ, Kwon J-W, La TY, Park CK, Choi JA. Characteristics of Retinal Nerve Fiber Layer Defect in Nonglaucomatous Eyes With Type II Diabetes. *Invest Ophthalmol Vis Sci.* 2016;57(10):4008-4015. doi:10.1167/iovs.16-19525.
26. Carpineto P, Toto L, Aloia R, et al. Neuroretinal alterations in the early stages of diabetic retinopathy in patients with type 2 diabetes mellitus. *Eye (Lond).* 2016;30(5):673-679. doi:10.1038/eye.2016.13.
27. El-Shazly AAE-F, Farweez YAT, Elewa LS, Elzankalony YA, Farweez BAT. Effect of Active and Passive Smoking on Retinal Nerve Fibre Layer and Ganglion Cell Complex. *J Ophthalmol.* 2017;2017:6354025. doi:10.1155/2017/6354025.
28. Dervişoğulları MS, Totan Y, Tenlik A, Yüce A, Güler E. Effect of smoking on retina nerve fiber layer and ganglion cell-inner plexiform layer complex. *Cutan Ocul Toxicol.* 2015;34(4):282-285. doi:10.3109/15569527.2014.975240.
29. Yoshida M, Take S, Ishikawa M, et al. Association of smoking with intraocular pressure in middle-aged and older Japanese residents. *Environ Health Prev Med.* 2014;19(2):100-107. doi:10.1007/s12199-013-0359-1.
30. Park H-YL, Park YG, Cho A-H, Park CK. Transneuronal retrograde degeneration of the retinal ganglion cells in patients with cerebral infarction. *Ophthalmology.* 2013;120(6):1292-1299. doi:10.1016/j.ophtha.2012.11.021.

31. Anjos R, Vieira L, Costa L, et al. Macular Ganglion Cell Layer and Peripapillary Retinal Nerve Fibre Layer Thickness in Patients with Unilateral Posterior Cerebral Artery Ischaemic Lesion: An Optical Coherence Tomography Study. *Neuroophthalmology*. 2016;40(1):8-15. doi:10.3109/01658107.2015.1122814.
32. He X-F, Liu Y-T, Peng C, Zhang F, Zhuang S, Zhang J-S. Optical coherence tomography assessed retinal nerve fiber layer thickness in patients with Alzheimer's disease: a meta-analysis. *Int J Ophthalmol*. 2012;5(3):401-405. doi:10.3980/j.issn.2222-3959.2012.03.30.
33. Kuang TM, Zhang C, Zangwill LM, Weinreb RN, Medeiros FA. Estimating Lead Time Gained by Optical Coherence Tomography in Detecting Glaucoma before Development of Visual Field Defects. *Ophthalmology*. 2015;122(10):2002-2009. doi:10.1016/j.ophtha.2015.06.015.

Figure legends

Figure 1a: Forest plots of meta-analyzed associations with pRNFLT from multivariable regression models (Age, sex, spherical equivalent, intraocular pressure and visual impairment). The beta-coefficients [95% Confidence Interval] show the influence of each parameter on pRNFLT within the respective study, the percentage represents the mathematically determined weighting of each study within the meta-analysis.

Figure 1b: Forest plots of meta-analyzed associations with pRNFLT from multivariable regression models (Smoking, hypertension, stroke and dementia). The beta-coefficients [95% Confidence Interval] show the influence of each parameter on pRNFLT within the respective study, the percentage represents the mathematically determined weighting of each study within the meta-analysis.

Table 1. Descriptive data for participating studies

Study	Years	City/country	pRNFLT measurements		N	Women (%)	Mean age in years ± SD	Mean global pRNFLT in microns ± SD
			OCT - Device	Details				
Alienor Study	2009-2011	Bordeaux, France	Spectralis OCT, Heidelberg Engineering	12° / 3.4mm diameter ring scan on ONH	529	62%	82.1±4.2	89.2±16.0
Coimbra Eye Study	2016-2017	Coimbra, Portugal			618	54%	71.8±6.2	96.8±12.0
Montrachet Study	2009-2013	Dijon, France			803	60%	82.0±3.7	90.3±13.7
LIFE Study	2011-2014	Leipzig, Germany			8351	52%	56.9±12.3	97.4±10.6
Rotterdam Study I	2009-2011	Rotterdam, the Netherlands	3D OCT 1000, Topcon Medical Systems	6.0 x 6.0mm on ONH	1287	57%	79.3±4.6	86.8±21.4
Rotterdam Study II	2011-2012		3D OCT 1000 and 2000, Topcon Medical Systems		1376	55%	72.4±4.9	98.2±17.2
Rotterdam Study III	2012-2013				2267	56%	62.2±5.6	104.7±12.5
Twins UK Study	2014-2016	UK (multiple cities)	iVue, Optovue	3.45 mm diameter ring scan on ONH	853	98%	61.8±12.2	96.4±9.8

pRNFLT=Peripapillary retinal nerve fiber layer thickness, SD=Standard deviation, OCT=Optical coherence tomography, ONH=Optic nerve head

Table 2a. Associations with peripapillary retinal nerve fiber layer thickness (pRNFLT) for each individual study

	Alienor (n=529)		Coimbra (n=618)		Montrachet (n=803)		LIFE (n=8351)	
	β (95% CI)	p	β (95% CI)	p	β (95% CI)	p	β (95% CI)	p
Age (per year)	-0.22 (-0.55; 0.11)	0.12	-0.09 (-0.24; 0.07)	0.28	-0.37 (-0.62; -0.12)	0.004	-0.10 (-0.12; -0.08)	<0.001
Female sex	3.98 (0.84; 7.11)	0.01	2.78 (0.70; 4.85)	0.01	3.85 (1.78; 5.92)	<0.001	1.36 (0.90; 1.82)	<0.001
BMI (per kg/m ²)	0.13 (-0.22; 0.48)	0.46	-0.10 (-0.30; 0.11)	0.37	0.33 (0.09; 0.57)	0.006	0.08 (0.03; 0.13)	0.002
SE (per diopter)	1.22 (0.50; 1.94)	<0.001	1.15 (0.60; 1.69)	<0.001	1.88 (1.44; 2.31)	<0.001	N/A	N/A
IOP (per mmHg)	-0.71 (-1.32; -0.10)	0.02	-0.52 (-0.84; -0.19)	0.002	-0.31 (-0.60; -0.03)	0.03	N/A	N/A
Visual impairment*	-6.42 (-12.47; -0.37)	0.04	-0.15 (-4.64; 4.34)	0.95	-0.49 (-4.46; 3.48)	0.81	N/A	N/A
Former smoker	-0.47 (-3.74; 2.81)	0.78	-0.57 (-3.67; 2.53)	0.72	2.39 (0.25; 4.53)	0.03	0.60 (0.06; 1.14)	0.03
Current smoker	1.48 (-4.99; 7.96)	0.65	1.51 (-5.74; 8.77)	0.68	1.58 (-4.33; 7.48)	0.60	1.43 (0.85; 2.02)	<0.001
Hypertension	1.33 (-1.40; 4.05)	0.34	-1.08 (-3.07; 0.91)	0.29	-0.62 (-2.55; 1.32)	0.53	-0.46 (-0.98; 0.07)	0.09
Diabetes	2.11 (-2.30; 6.52)	0.35	1.34 (-0.88; 3.55)	0.24	-2.08 (-5.29; 1.13)	0.21	-1.25 (-2.03; -0.46)	0.002
Stroke	-3.53 (-14.05; 6.99)	0.51	-2.85 (-7.60; 1.91)	0.24	0.02 (-5.20; 5.25)	0.99	-2.96 (4.56; -1.36)	<0.001
Dementia	-5.11 (-12.06; 1.84)	0.15	N/A	N/A	N/A	N/A	N/A	N/A

Results from the multivariable regression models; N/A= not available

*Visual impairment as defined by the World Health Organization (<0.3 decimal BCVA); BMI=Body Mass Index, SE=Spherical equivalent, IOP=Intraocular pressure

Table 2b. Associations with peripapillary retinal nerve fiber layer thickness (pRNFLT) for each individual study (continued)

	Rotterdam 1 (n=1287)		Rotterdam 2 (n=1376)		Rotterdam 3 (n=2267)		Twins UK (n=853)	
	β (95% CI)	p	β (95% CI)	p	β (95% CI)	p	β (95% CI)	p
Age (per year)	-0.61 (-0.87; -0.36)	<0.001	-0.91 (-1.08; -0.73)	<0.001	-0.50 (-0.59; -0.41)	<0.001	-0.24 (-0.29; -0.18)	<0.001
Female sex	3.70 (1.23; 6.16)	0.003	1.36 (-0.31; 3.03)	0.11	0.49 (-0.50; 1.48)	0.33	3.58 (-1.01; 8.16)	0.13
BMI (per kg/m ²)	0.11 (-0.18; 0.40)	0.46	0.00 (-0.002; 0.002)	1.00	0.12 (0.00; 0.24)	0.05	-0.07 (-0.19; 0.05)	0.27
SE (per diopter)	1.09 (0.58; 1.60)	<0.001	1.67 (1.32; 2.02)	<0.001	1.29 (1.10; 1.49)	<0.001	1.28 (1.02; 1.55)	<0.001
IOP (per mmHg)	-0.49 (-0.84; -0.14)	0.005	-0.65 (-0.91; -0.39)	<0.001	-0.05 (-0.23; 0.12)	0.56	-0.05 (-0.26; 0.16)	0.62
Visual impairment*	-11.27 (-17.93; -4.61)	<0.001	-7.68 (-13.80; -1.55)	0.01	-9.56 (-14.87; -4.24)	<0.001	N/A	N/A
Former smoker	2.60 (-0.05; 5.25)	0.06	0.87 (-0.95; 2.68)	0.36	-0.13 (-1.20; 0.95)	0.82	N/A	N/A
Current smoker	2.96 (-1.48; 7.40)	0.19	3.86 (0.94; 6.78)	0.009	1.34 (-0.18; 2.87)	0.08	N/A	N/A
Hypertension	2.08 (-1.71; 5.86)	0.28	0.46 (-1.86; 2.77)	0.70	-1.37 (-2.43; -0.31)	0.01	N/A	N/A
Diabetes	-0.54 (-3.45; 2.36)	0.71	-1.72 (-3.92; 0.48)	0.12	0.40 (-1.30; 2.10)	0.65	-4.29 (-8.34; -0.24)	0.04
Stroke	1.25 (-2.83; 5.32)	0.55	-2.06 (-5.67; 1.55)	0.26	-1.22 (-4.19; 1.76)	0.42	-1.51 (-5.49; 2.48)	0.46
Dementia	-4.17 (-8.82; 0.48)	0.08	-0.55 (-5.96; 4.86)	0.84	-1.27 (-24.41; 21.86)	0.91	N/A	N/A

Results from the multivariable regression models; N/A= not available

*Visual impairment as defined by the World Health Organization (<0.3 decimal BCVA); BMI=Body Mass Index, SE=Spherical equivalent, IOP=Intraocular pressure

Table 3. Meta-analyzed associations with peripapillary retinal nerve fiber layer thickness (pRNFLT)

	All participants			Excluding known glaucoma		
	β (95% CI)	p	I^2	β (95% CI)	p	I^2
Age (per year)	-0.38 (-0.57; -0.18)	<0.001	97%	-0.35 (-0.60; -0.10)	0.006	97%
Female sex	2.17 (1.15; 3.20)	<0.001	69%	1.79 (0.93; 2.65)	<0.001	59%
BMI (per kg/m ²)	0.06 (-0.02; 0.14)	0.15	54%	0.09 (0.00; 0.18)	0.05	53%
SE (per diopter)	1.39 (1.19; 1.59)	<0.001	49%	1.36 (1.16; 1.57)	<0.001	40%
IOP (per mmHg)	-0.36 (-0.56; -0.15)	<0.001	74%	-0.42 (-0.65; -0.20)	<0.001	71%
Visual impairment*	-5.50 (-9.37; -1.64)	0.005	69%	-4.75 (-9.12; -0.38)	0.03	77%
Former smoker	0.58 (0.14; 1.02)	0.009	0%	0.79 (-0.01; 1.60)	0.05	43%
Current smoker	1.53 (1.00; 2.06)	<0.001	0%	1.49 (0.97; 2.02)	<0.001	0%
Hypertension	-0.54 (-1.01; -0.07)	0.03	4%	-0.62 (-1.11; -0.13)	0.01	7%
Diabetes	-0.69 (-1.69; 0.31)	0.18	40%	-0.33 (-1.32; 0.67)	0.52	40%
Stroke	-1.94 (-3.17; -0.72)	0.002	7%	-2.06 (-3.27; -0.84)	<0.001	2%
Dementia	-3.11 (-6.22; 0.01)	0.05	0%	-2.66 (-5.86; 0.55)	0.10	0%

All participants – results of the meta analysis of the multivariable regression models including all participants (n=16084)

Excluding known glaucoma – results of the meta analysis of the multivariable regression models excluding the 619 known glaucoma patients (n=14695, without TwinsUK Study data, since no data on glaucoma were available); I^2 =Heterogeneity of covariate in the meta-analysis; *Visual impairment as defined by the World Health Organization (<0.3 decimal BCVA); BMI=Body Mass Index, SE=Spherical equivalent, IOP=Intraocular

pressure
